

Inequalities in the Management of Diabetic Kidney Disease in UK Primary Care: A Cross-Sectional Analysis of A Large Primary Care Database

Running title: Inequalities in the management of DKD

Authors

Katherine Phillips¹, Jonathan M Hazlehurst^{1,2,3}, Christelle Sheppard⁴, Srikanth Bellary^{3,5}, Wasim Hanif^{2,3}, Muhammad Ali Karamat^{2,3}, Francesca L Crowe¹, Anna Stone⁶, G Neil Thomas¹, Javeria Peracha⁷, Anthony Fenton⁸, Christopher Sainsbury^{1,9}, Krishnarajah Nirantharakumar^{1,10}, Indranil Dasgupta^{11,12}

Affiliations

¹ Institute of Applied Health Research, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK.

² Centre for Endocrinology, Diabetes and Metabolism, Birmingham Health Partners, Birmingham, UK.

³ Department of Diabetes and Endocrinology, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK.

⁴ Sandwell and West Birmingham Hospital NHS Trust, Birmingham, UK.

⁵ School of Health and Life Sciences, Aston University, Birmingham, UK.

⁶ Royal Wolverhampton NHS Trust, UK

⁷ Renal Unit, Royal Wolverhampton NHS Trust, UK

⁸ Department of Renal Medicine, Royal Stoke University Hospital, University Hospitals of North Midlands NHS Trust

⁹ Department of Diabetes, Gartnavel General Hospital, Glasgow, UK.

¹⁰ Midlands Health Data Research UK, University of Birmingham, Birmingham, UK.

¹¹ Department of Renal Medicine, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK.

¹² Warwick Medical School, University of Warwick, UK.

Corresponding author

Indranil Dasgupta Indranil.Dasgupta@uhb.nhs.uk

Word count (excluding abstract): 4042 Abstract word count: 288 Tables: 1

Conflict of interest disclosures: ID received research grants from Sanofi and Medtronic and attended advisory board and delivered talks for GSK, Vifor, Sanofi and AstraZeneca. MAK has received speaker honoraria and/or advisory board fees from Novo Nordisk, Eli Lilly, B.I, Astrazeneca, Sanofi and Abbott Diabetes care.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1111/dme.15153](https://doi.org/10.1111/dme.15153)

Novelty statement

What is already known?

- There are inequalities in outcomes for patients with diabetes and CKD between different socioeconomic groups and ethnicities.

What this study found

- Women with DKD were less likely to be appropriately monitored, prescribed appropriate medications or treated to target.
- People from socioeconomically deprived backgrounds were less likely to be appropriately monitored, prescribed appropriate medications or treated to target.
- People of black ethnicity were less likely to be prescribed statins than white ethnic group.

What are the implications of the study?

- Further research is needed to test low-cost interventions that improve and standardise the management of DKD in primary care.

Acknowledgements

Funding and assistance

This project whilst not specifically funded was supported by funding to the STOP Diabetes project team by the NIHR Clinical Research Network West Midlands.

Author contribution and guarantor statement

All authors were involved in the conception and design of this work. KP conducted the analysis of data. KP, JH, CS and ID drafted the work and all authors edited, reviewed and approved the final version of the manuscript.

Data and resource availability statement

The data that support the findings of this study are available from IQVIA Medical Research Data but restrictions apply to the availability of these data, which were used under license for the current study and therefore are not publicly available. Data are however available from the authors upon reasonable request and with permission of IQVIA Medical Research Data.

Ethics approval

Scientific review committee approval for this analysis was obtained from IMRD in October 2021 (SRC reference 21SRC051).

Abstract

Objective: To determine differences in the management of Diabetic Kidney Disease (DKD) relevant to patient sex, ethnicity and socioeconomic group in UK primary care.

Research Design and Methods: A cross-sectional analysis as of 1st January 2019 was undertaken using IQVIA Medical Research Data (IMRD) dataset, to determine the proportion of people with DKD managed in accordance with national guidelines, stratified by demographics. Robust Poisson regression models were used to calculate adjusted risk ratios (aRR) adjusting for age, sex, ethnicity, and social deprivation.

Results: Of the 2.3 million participants in the database, 161,278 had type 1 or 2 diabetes of whom 32,905 had DKD.

Compared to men, women were less likely to have creatinine: aRR 0.99 (95% CI 0.98-0.99), albumin creatine ratio (ACR): aRR 0.94 (0.92-0.96), blood pressure (BP): aRR 0.98 (0.97-0.99), glycosylated haemoglobin (HbA1c): aRR 0.99 (0.98-0.99), and serum total cholesterol: aRR 0.97 (0.96-0.98) measured; achieve BP (<140/80 mmHg): aRR 0.95 (0.94-0.98) or total cholesterol (< 5 mmol/L) targets: aRR 0.86 (0.84-0.87); or be prescribed renin-angiotensin-aldosterone system (RAAS) inhibitors: aRR 0.92 (0.90-0.94) or statins: aRR 0.94 (0.92-0.95) in the previous year.

Compared to the least deprived areas, people from the most deprived areas were less likely to have BP measurements: aRR 0.98 (0.96-0.99); achieve BP (<140/80 mmHg): aRR 0.91 (0.8-0.95) or HbA1c (< 58 mmol/mol): aRR 0.88 (0.85-0.92) targets, or be prescribed RAAS inhibitors: aRR 0.91 (0.87-0.95), but were 8% more likely to be prescribed a statin: aRR 1.08 (1.06-1.11). Compared to patients of white ethnicity; patients of black ethnicity were less likely to be prescribed statins aRR 0.91 (0.85-0.97).

Conclusions: There are unmet needs and inequalities in the management of DKD in the UK. Addressing these could reduce the increasing human and societal cost of managing DKD.

Keywords: DKD, inequality, diabetes, ethnicity

Introduction

Diabetic kidney disease (DKD) is a chronic complication of diabetes and affects 30-40% of people with type 2 diabetes.(1) It accounts for around 42% cases of chronic kidney disease (CKD) globally.(2) DKD is an established risk factor for all-cause and cardiovascular mortality and end stage kidney disease (ESKD); 30.2% of patients requiring dialysis in the United Kingdom (UK) have DKD as the primary kidney disease. (3)

Chronic kidney disease (CKD) and Type 2 diabetes are more prevalent in people from lower socioeconomic groups and those of black and South Asian ethnicities.(4) People from lower socioeconomic groups are more likely to develop DKD,(1) and those from black and South Asian ethnicities with DKD are more likely to progress to more advanced stages of CKD, and along with those socioeconomically deprived, are more likely to start dialysis. CKD is more prevalent amongst women, but men are more likely to receive dialysis.(4)

Early recognition and treatment of DKD are important in order to modify cardiovascular disease risk and to limit progression of disease to End Stage Kidney Disease (ESKD). In the UK, clinical practice guidelines exist, providing evidence-based recommendations for the management of DKD. These include lifestyle advice, pharmacotherapy for hypertension, lipids and glycaemic control, and recommendations for disease monitoring and treatment targets.(5-7) The NICE recommendations for both CKD and diabetes (8) (9) are aimed at the primary care management of both conditions in England and have been updated in 2021 to include more recent evidence.

Data from the National Diabetes Audit in England and Wales showed that people with diabetes who have fewer care processes had a higher mortality rate. There was a significant association between the number of care processes completed and ethnicity, but not deprivation, and the association of completion of care processes and mortality also differed significantly by ethnicity but not deprivation. However, data on the individual targets that were achieved in these different at-risk groups were not available (10) thus making it difficult to examine the full extent of variation in care and outcomes by ethnicity or deprivation.

In view of the relationship between ethnicity and deprivation and CKD management in the UK, as well as the association of T2DM and progression to ESKD with deprivation and certain ethnic groups, we set out to examine this further using a large, comprehensive primary care dataset from 700 general practices across the UK. The aim of this study was to investigate whether there were any significant variations in the management of DKD in the UK primary care in relation to sex, ethnicity and socioeconomic status.

Methods

Study design and data source

We performed a cross-sectional analysis of the IQVIA Medical Research Data (IMRD) database using an index date of 1st January 2019, to determine the proportion of patients with DKD managed in accordance with a range of care quality metrics derived from the NICE Diabetes and CKD guidelines and guidelines from the Association of British Clinical Diabetologists (ABCD) (5-9, 11).

IMRD is an anonymised database of electronic primary care records from general practices in the UK using Vision software. All participating practices contribute coded data on patient characteristics, prescriptions, consultations, diagnoses and primary care investigations. IMRD contains records from over 700 general practices spread across all four nations of the UK, with more than 15 million patient records (22% of the UK population). Around 3.7 million of these are active at any given time point. IMRD is largely representative of the UK population in terms of demographics and morbidity prevalence.(12) Symptoms, clinical examinations, and diagnoses in IMRD are recorded using a hierarchical clinical coding system called Read codes.(13) Practices were eligible for inclusion in the study from 12 months after the latest of either practice acceptable mortality recording date (AMR) (a standard measure of sufficient data quality) (14) or Vision software installation date.

DKD was defined as the presence of Read codes for both diabetes and CKD. The data extraction and cohort selection were facilitated using the data extraction for epidemiological research (DExtER) tool.(15)

Study population

People aged 18 years and older who had been contributing data to the IMRD for at least 12 months as of the 1st of January 2019 were eligible for this analysis. From this, only people with a diagnosis of type 1 or type 2 diabetes and CKD were included.

Definition of variables

Diagnoses of CKD (stages 1-5) and type 1 or type 2 diabetes were defined as the presence of a Read Code for these conditions that were recorded before the 1st of January 2019. The most recent record of sex, ethnicity, Townsend deprivation score (based on the patients' postcode), and body mass index (BMI) before the 1st of January 2019 were included in the dataset. BMI was categorised using the World Health Organization classification.

A record of prescription for statins and renin-angiotensin-aldosterone system (RAAS) inhibitors within the past year was used to determine current treatment. Prescription of RAAS inhibitors was considered appropriate if the patient had an albumin creatinine ratio (ACR) ≥ 3 mg/mmol (8). Statins were considered appropriate for all patients in this analysis as NICE guidance advises offering statins to all patients with CKD. (16)

Key outcome measures (6) (7) (8) (9)

1. A record for blood pressure (BP), cholesterol, creatinine (to calculate estimated glomerular filtration rate; eGFR), and albumin creatine ratio (ACR) measurements within the previous year.
2. Appropriate prescription of statins and RAAS inhibitors as defined above.
3. BP maintained at or below target ($\leq 140/80$ mmHg)
4. Glycosylated haemoglobin (HbA1c) maintained at or below target (≤ 58 mmol/mol (7.5%))
5. Serum total cholesterol at or below target (≤ 5 mmol/L)

For all these key outcome measures, absence of data for that variable was assumed to mean that the monitoring has not been carried out or the medication had not been prescribed as is the nature with this type of data.

Statistical analysis

Age at the study entry was calculated and categorised into the following groups: 18–30, 31–40, 41–50, 51–60, 61–70, 71–80 (reference) and ≥ 81 years. Ethnicity was categorised into five groups: white (reference), black, South Asian, mixed ethnicity, and other. Socioeconomic group was based on the Townsend deprivation score and divided into quintiles with the lowest (reference) corresponding to the least deprived and the highest to the most deprived.

Characteristics of participants were reported using appropriate descriptive statistics (mean for normally distributed continuous variables, median and interquartile range (IQR) for non-normally distributed variables, and proportions for categorical variables). For each of the key outcome measures, the proportion of people who received appropriate management within the previous year was calculated overall and stratified by age, sex, ethnicity, and socioeconomic status. Robust Poisson regression models using a log-link function were used to estimate risk ratios for the key outcome measures among those with DKD according to age, sex, ethnicity, and Townsend deprivation score and for the adjusted risk ratios (aRR) and 95% confidence intervals (CI) after adjusting for age, sex, ethnicity, and Townsend deprivation score (where appropriate). To account for similarity of outcomes within GP practices, statistical inference from the log binomial regression model was based on cluster-robust standard errors. Data were missing for 27% for socioeconomic group, and 47% of patients for ethnicity. To ensure that the same patients were being compared in all analyses those with a missing value for each particular variable were assigned to a separate category for that variable and included in the regression analysis.

All statistical analyses were performed using Stata statistical software, V.15.1 (StataCorp, College Station, Texas, USA). Two-sided p values < 0.05 were considered to be statistically significant.

Results

The records of 2,293,244 patients 18 years and older were eligible for inclusion in the analysis, of whom 161,278 (7.03%) had a code for type 1 (14,596) or type 2 diabetes (146,682). Of those with diabetes, 32,556 (20.2%) also had a code for CKD and were included in the analysis.

Baseline characteristics are presented in **Table 1**. Men represented 48.7% of the cohort, the mean age was 76.7 years. Of the 32,556 people with DKD, 1,987 (6.1%) had a diagnosis of type 1 diabetes and 30,579 (93.9%) had a diagnosis of type 2 diabetes.

Renal function

Of the 32,556 with DKD, 30,171 (92.7%) had their creatinine and 19,433 (59.7%) had ACR measured within the previous year (**Figure 1 and Supplementary Table 1**). Compared to those age 71-80 years, people aged 81 years and older were 9% less likely to have ACR measured: aRR 0.91 (95% CI 0.89-0.93) and people aged 18-30 years were 69% less likely to have ACR measured (95% CI 0.11-0.86). Compared to men, women had a significantly lower risk of having their serum creatinine: aRR 0.99 (95% CI 0.98-0.99) and ACR: aRR 0.94 (95% CI 0.92-0.96) measured within one year (**Figure 2 and Supplementary Table 2**).

Blood pressure

Overall, 30,202 (92.7%) of patients with DKD had their BP measured within a year. However, only 20,915 (64%) had BP at or below the target of 140/80 mmHg (Figure 1 and Supplementary Table 1). Results in Figures 2 & 3 and Supplementary Table 2 show that those age 81 years and older were significantly less likely to have their blood pressure measured in the last year (aRR 0.98 (95% CI 0.97-0.99) and for it to be below target: aRR 0.96 (0.95-0.98) and as were those aged 31-40 years (blood pressure measured: aRR 0.9 (0.83-0.97) below target: aRR 0.71 (0.6-0.83)). Women were at 2% less likely to have their BP measured within a year: aRR 0.98 (95% CI 0.98-0.99) and have a BP below target: aRR 0.95 (0.94-0.97). People living in the most deprived areas were 2% less likely to have their BP measured within a year: aRR 0.98 (95% CI 0.96-0.99) and 9% less likely for BP to be on target: aRR 0.91 (0.87-0.95) compared to those in the least deprived areas.

HbA1c

There were 29,444 (90.4%) patients with DKD who had their HbA1c measured within one year (Figure 1 and supplementary Table 1). However, only 18,876 (58.0%) had their HbA1c at the target of 58 mmol/mol or lower (Figure 1 and Supplementary Table 1). Those age 81 years and older were 3% less likely to have their HbA1c measured: aRR 0.97 (95% CI 0.96-0.98), but of those that did, this older age group were slightly more likely to have HbA1c lower than 58 mmol/mol: aRR 1.04 (95% CI 1.02-1.06) compared to those age 71-80 years. Those aged 18-30 were also less likely to have had their HbA1c below target (aRR 0.33 (0.12-0.94)). Women were less likely to have their HbA1c measured within the past year: aRR 0.99 (95% CI 0.98-0.99). People living in the most deprived areas were 12% less likely to have their HbA1c measurement lower than 58 mmol/mol (7.5%): aRR 0.88 (95% CI 0.85- 0.92) (Figures 2 & 3 and Supplementary Table 2).

Serum total cholesterol

In the cohort overall, 26,843 (82.8%) of patients had their serum cholesterol measured within a year and total cholesterol was ≤ 5 mmol/L for 22,655 (69.6%) of patients (Figure 1 and supplementary Table 1). Those age 81 years and older were 6% less likely to have their serum cholesterol measured: aRR

0.94 (95% CI 0.93-0.95) and were also 7% less likely to meet the target of ≤ 5 mmol/L: aRR 0.93 (95% CI 0.91-0.94). Those aged 31-40 were 23% less likely to have their cholesterol measured (95% CI 0.68-0.86) and 39% less likely for it to be on target (95% CI 0.5-0.75). Women were less likely to have their serum cholesterol measured within a year: aRR 0.97 (0.96-0.98) and 14% less likely for their total cholesterol to be ≤ 5 mmol/L (aRR 0.86 (0.84-0.87)). People of South Asian ethnicity were 9% more likely to have their serum cholesterol on target: aRR 1.09 (1.03-1.15) (Figures 2 & 3 and Supplementary Table 2).

RAAS inhibitors

RAAS inhibitors were indicated in 18,901 of the study cohort. 12,842 (68.4%) of these were prescribed RAAS inhibitors within the previous year (Figure 1 and Supplementary Table 1). Results in Figure 4 and Supplementary Table 2 show that those age 81 years and older were 13% less likely to be prescribed a RAAS inhibitor (aRR 0.87 (95% CI 0.85-0.89)). Those aged 31-40 were 33% less likely to be prescribed RAAS inhibitors (95% CI 0.55-0.81). Women were 8% less likely to have been appropriately treated with RAAS inhibitors within the previous year: aRR 0.92 (0.90-0.94). People living in the most deprived areas were 9% less likely to have been prescribed RAAS inhibitors: aRR 0.91 (95% CI 0.87-0.95).

Statins

In the DKD cohort, 25,374 (77.9%) were prescribed statins within the previous year (Figure 1 and Supplementary Table 1). Those 81 years were 11% less likely to be prescribed a statin: aRR 0.89 (95% CI 0.88-0.90) and those aged 31-40 were 61% less likely to be prescribed a statin (95% CI 0.31-0.5). Women were significantly less likely to have been prescribed statins within the previous year: aRR 0.94 (95% CI 0.92-0.95) (Figure 1). People of black ethnicity were 9% less likely to have been prescribed statins within the previous year: aRR 0.91 (95% CI 0.85-0.97), and people of South Asian ethnicity were 7% more likely to have been prescribed statins within the previous year: aRR 1.07 (1.04-1.11). People living in the most deprived areas were also 8% more likely to have been prescribed statins within the previous year: aRR 1.08 (1.06-1.11) (Figure 4 and Supplementary Table 2).

Including an adjustment for the type of diabetes made little difference to the risk estimates of the key outcome measures for sex, ethnicity and social deprivation. However, patients aged 18-30 were no longer at significant risk of having ACR measured or HbA1c being on target when type of diabetes was included as an adjustment (supplementary table 3).

Discussion

This analysis of the management of DKD in primary care demonstrates areas of significant unmet need and inequality.

Overall management

Over 90% of patients with DKD in this study had HbA1c, BP, and serum creatinine measured within the previous year. This is consistent with results from the National Diabetes Audit (NDA) on people with type 2 diabetes in England and Wales for 2018-19 (17). Fewer patients in this study had serum cholesterol measured (83.7%) and slightly higher percentage had ACR measured (58%) than in the NDA analysis (97.8% and 49.3% respectively). However, with regard to meeting the BP target of $\leq 140/80$ mmHg (the target used in the NDA), a difference was observed between the two analyses (75% in the NDA v 64.2% in this study). (17) Our results are similar to a previous analysis of NDA data from 2007-08 of patients with DKD where 62% of patients were below the BP target (18). In the present work 58% of patients were at or below the HbA1c target (58 mmol/mol) which is in contrast to previous studies where 32% of those with type 1 and 68% of those with type 2 met this glycaemic target (18). It is interesting to note that there has been very little change in the achievement of either BP or glycaemic target in people with DKD in the UK in over a decade.

Following the introduction of financial incentives to monitor urinary markers of CKD through the Quality Outcomes Framework (QOF) in 2010/11, improvements in the rates of annual ACR testing were reported. (19) However, this particular QOF indicator was “retired” in 2014, perhaps contributing to the overall lower rates of testing found in this study and in the recent NDA report. (20)

We also observed that older adults (81+ years) tended to have fewer care processes measured and less likely to meet the treatments targets compared to the younger age groups. This could possibly be due to difficulties in attending appointments and more relaxed treatment targets due to advancing age and frailty.

Management disparity by age

A U-shaped relationship was seen between age and many of the key outcome measures. Compared to those ages 71-80, patients aged over 81 years were less likely to have had ACR, BP, HbA1c and cholesterol measured BP and cholesterol on target and RAAS inhibitors and statins prescribed. Patients aged 31-40 were significantly less likely to have had any of the key outcomes measured. Patients aged 18-30 were less likely to have had ACR measured, HbA1c on target or statins prescribed (although including an adjustment term for type of diabetes meant that there was no longer a significant association for ACR measurement or achievement of HbA1c targets). Low numbers of patients in this age group may account for the lack of significance for some of the results.

A similar U-shaped relationship with age was found in Italian cohort study which analysed compliance to care processes for the management of diabetes, congestive heart failure and coronary heart disease. Young adults and those over 75 had lower levels of achievement for almost all quality indicators examined than those aged 65-74 (21). A London study also examined the impact of age on achievement of quality indicators in diabetes and found those aged 18-44 years were less likely to have creatinine, ACR, blood pressure, cholesterol, and HbA1c measured compared to the older age groups (22).

Potential reasons for those aged 65-74 years having better achievement of key outcomes include ease of access (opening hours of GP practices may mean they are more accessible to those who are retired) and better adherence to management (21, 22). In older and, potentially, more frail adults, it may be

appropriate to adjust treatment targets, which could account for the reduced likelihood of being managed in accordance with guidance in those over 81 years.

Management disparity by sex

Compared to men, women with DKD were less likely to have biochemical markers measured within the previous year, have their BP and cholesterol measurements below or at the target, or were prescribed RAAS or statins. Although some of the differences we found were small, inequalities by sex in the management of cardiovascular risk factors have been described previously. Lower prescribing rates for statins and RAAS inhibitors in women have been reported in a systematic review (23). Moreover, results from a multi-country study showed that women with established coronary heart disease were less likely than men to achieve all treatment targets including glucose and cholesterol. However, women were more likely to meet BP targets, which is in contrast to the current study.(24) The finding that women were less likely to be managed to treatment targets is well established in the secondary prevention of cardiovascular disease (25) (26). There are several potential reasons for these differences including reduced prescription of medicines for secondary prevention (27), underestimation of cardiovascular risk of women by clinicians (28), and a lower awareness of cardiovascular risk by women (29) despite better engagement with and attendance at follow up (30). The prior finding of reduced appropriate prescribing for women(27) is consistent with results from this study of lower prescribing of RAAS and statins. These gender disparities for management of cardiovascular disease are similar to that within the management and attainment of treatment targets for type 2 diabetes (31).

Management disparity by ethnicity

There were some inequalities noted between ethnic groups. Compared to white ethnicity, people of black ethnicity were less likely to be prescribed statins. Interestingly, people of South Asian ethnicity were more likely to be prescribed statins than people of white ethnicity and meet the target for serum total cholesterol < 5mmol/L. Disparities in the management of DKD have been described between ethnic groups in the UK previously. People of black ethnicity were less likely to have their BP controlled to below target in a cross-sectional study of patients in London (32). The relatively small numbers of patients in some of the ethnicity categories may mean that our study was not powered to detect differences between groups and almost half of the study population did not have their ethnicity recorded, which means some of these findings should be interpreted in light of this. It would be imperative to improve recording of ethnicity in primary care records, as the first step, to address the disparities in the management noted in this study.

In England, Wales and Northern Ireland, QOF incentivises maintaining a register of patients with CKD stages 3-5 and diabetes registers that include aspects of diabetes care (16). Despite this, results from the NDA show that approximately 30% of patients meeting biochemical criteria for CKD in primary care do not have a diagnostic code for CKD (17). Lower levels of CKD diagnoses are associated with greater deprivation and being in a non-white ethnicity group. Being non-coded for CKD is also associated with poorer management (33) (but these people would not have been captured in this analysis). It may be the case that, once identified as having DKD, people from certain ethnic groups are generally managed in accordance with guidance. This would be in keeping with results of the National Diabetes Audit (17).

Management disparity by deprivation

People from more deprived areas were less likely to have BP measured; less likely to have their HbA1c or BP below the target, and less likely to be prescribed RAAS inhibitors but more likely to be prescribed statins compared with those from less deprived areas. Other studies have reported suboptimal diabetes control in patients of lower socioeconomic status and that high social deprivation was independently associated with worse control (34). Several reasons for this association have been suggested. Health literacy, a person's capacity to access, understand, appraise, and apply health information, has been found to be lower in people of lower socioeconomic status. Increasing medical complexity, reduced capacity to cope and reduced access to care, as well as being more likely to serially miss appointments are potential factors contributing to poorer health outcomes in lower socioeconomic groups (4, 35, 36).

Strengths and Limitations

The cohort of more than 30,000 people with DKD comes from a large primary care dataset which has been demonstrated to be representative of the UK population (23). Therefore, the results are generalisable to the UK primary care population. To our knowledge, this is the first UK study looking specifically at the management of DKD patients in relation to sex, social deprivation, and ethnicity.

However, there are several limitations to this study. This is a cross-sectional analysis and therefore management over time was not captured. The time period of the study was prior to the COVID-19 pandemic, so this may not reflect the current state of management of these patients. The prevalence of CKD in this study is lower than would be expected in a diabetic cohort, likely due to poor coding. Almost half (47%) of patients did not have ethnicity coded and a measure of deprivation was missing for 27% and therefore some inequalities between ethnic groups and deprivation may not have been detected. The study only considered management within general practice, as we were not able to link IMRD (primary care) to secondary care datasets. Therefore, patients managed within secondary care may have been misclassified as not receiving appropriate management.

For our analysis, we employed standardised cut-offs for BP and HbA1c supported by previously published guidance (9-13,16). These may not be appropriate for patients who had had previous drug reactions, contra-indications, or individualised targets (for example those with frailty or on dialysis). Since the time period considered in this study, NICE has recommended to consider starting sodium-glucose co-transporter-2 (SGLT2) inhibitors in people with DKD and proteinuria. (13) As there is evidence of inequalities in the prescription of these drugs for diabetes, it would be worth considering prescription of SGLT2 inhibitors in future studies (29).

Conclusions

This cross-sectional analysis of a large primary care database identifies significant inequalities in the management of DKD in the UK. Improving the awareness of heart and kidney disease among people living with diabetes, engaging them more in their own care and the development of low-cost interventions targeting the improvement of management of DKD in primary care are required to mitigate the variation in care identified. Given the difficulties in addressing several risk factors in a limited healthcare setting low-cost multicomponent strategies aimed at risk factor identification and management (37) will be central to the future of diabetes care to minimise the future risk of complications. An implementation trial, or a hybrid implementation-effectiveness trial, of multi-component interventions based on current national guidelines may help to facilitate embedding of such interventions within primary care, with a view to improving access to treatment in at-risk patients, which is crucial in the context of an increasing prevalence of DKD nationally.

Figure Legends

Figure 1: The proportion of patients appropriately monitored and managed in accordance with guidelines stratified by sex, ethnicity and Townsend deprivation score

Figure 2: Adjusted risk ratios for the monitoring of creatinine, 2A ACR, 2B BP, 2C cholesterol, 2D creatinine and 2E HbA1c between age categories, sex, ethnicities, and Townsend deprivation score

Figure 3: Adjusted risk ratios for the achievement of 3A BP, 3B cholesterol and 3C HbA1c targets between age categories, sex, ethnicities, and Townsend deprivation score

Figure 4: Adjusted risk ratios for the appropriate prescribing of 4A RAAS inhibitors and 4B statins between age categories, sex, ethnicities, and Townsend deprivation score

References

1. Retnakaran R, Cull CA, Thorne KI, Adler AI, Holman RR. Risk factors for renal dysfunction in type 2 diabetes: U.K. Prospective Diabetes Study 74. *Diabetes*. 2006;55(6):1832-9.
2. Xie Y, Bowe B, Mokdad AH, Xian H, Yan Y, Li T, et al. Analysis of the Global Burden of Disease study highlights the global, regional, and national trends of chronic kidney disease epidemiology from 1990 to 2016. *Kidney Int*. 2018;94(3):567-81.
3. UK Renal Registry (2022) UK Renal Registry 24th Annual Report – data to 31/12/2020, Bristol, UK.

Available from <https://ukkidney.org/audit-research/annual-report>.

4. Kidney Research UK. Kidney health inequalities in the United Kingdom. 2018. p. 85-90.
5. Dasgupta I, Banerjee D, Chowdhury TA, De P, Wahba M, Bain S, et al. Association of British Clinical Diabetologists (ABCD) and Renal Association clinical guidelines: Hypertension management and renin-angiotensin-aldosterone system blockade in patients with diabetes, nephropathy and/or chronic kidney disease. *British Journal of Diabetes*. 2017;17(4):160-4.
6. Mark PB, Winocour P, Day C. Management of lipids in adults with diabetes mellitus and nephropathy and/or chronic kidney disease: summary of joint guidance from the Association of British Clinical Diabetologists (ABCD) and the Renal Association (RA). *British Journal of Diabetes*. 2017;17(2):64-72.
7. Winocour P, Bain SC, Chowdhury TA, De P, Pokrajac A, Fogarty D, et al. Managing hyperglycaemia in patients with diabetes and diabetic nephropathy—chronic kidney disease Summary of recommendations 2018. *British Journal of Diabetes*. 2018;18(2):78-91.
8. The National Institute for Health and Care Excellence. Chronic kidney disease in adults: assessment and management. Clinical Guideline 182. London: NICE 2015.
9. The National Institute for Health and Care Excellence. Type 2 Diabetes in adults: management. NICE Guideline 28. London: NICE 2015.
10. Holman N, Knighton P, O'Keefe J, Wild SH, Brewster S, Price H, et al. Completion of annual diabetes care processes and mortality: A cohort study using the National Diabetes Audit for England and Wales. *Diabetes Obes Metab*. 2021;23(12):2728-40.
11. NHS Digital. Quality and Outcomes Framework (QOF) business rules v42 2020-2021 baseline release. London: NHS Digital, 2020.
12. Blak BT, Thompson M, Dattani H, Bourke A. Generalisability of The Health Improvement Network (THIN) database: demographics, chronic disease prevalence and mortality rates. *Inform Prim Care*. 2011;19(4):251-5.
13. Booth N. What are the Read Codes? *Health Libr Rev*. 1994;11(3):177-82.
14. Maguire A, Blak BT, Thompson M. The importance of defining periods of complete mortality reporting for research using automated data from primary care. *Pharmacoepidemiol Drug Saf*. 2009;18(1):76-83.
15. Gokhale KM, Chandan JS, Toulis K, Gkoutos G, Tino P, Nirantharakumar K. Data extraction for epidemiological research (DExtER): a novel tool for automated clinical epidemiology studies. *Eur J Epidemiol*. 2021;36(2):165-78.
16. The National Institute for Health and Care Excellence. Cardiovascular disease: risk assessment and reduction, including lipid modification. Clinical Guideline 181. London: NICE 2014.
17. NHS Digital. Report 1: Care Processes and Treatment Targets, 2018-2019. National Diabetes Audit. 2020.
18. Hill CJ, Cardwell CR, Patterson CC, Maxwell AP, Magee GM, Young RJ, et al. Chronic kidney disease and diabetes in the national health service: a cross-sectional survey of the U.K. national diabetes audit. *Diabetic medicine : a journal of the British Diabetic Association*. 2014;31(4):448-54.

19. Feakins B, Oke J, McFadden E, Aronson J, Lasserson D, O'Callaghan C, et al. Trends in kidney function testing in UK primary care since the introduction of the quality and outcomes framework: a retrospective cohort study using CPRD. *BMJ Open*. 2019;9(6):e028062.
20. NHS Employers. 2015/16 General Medical Services (GMS) contract Quality and Outcomes Framework (QOF). 2015. p. 159.
21. Buja A, Damiani G, Gini R, Visca M, Federico B, Donato D, et al. Systematic Age-Related Differences in Chronic Disease Management in a Population-Based Cohort Study: A New Paradigm of Primary Care Is Required. *PLoS ONE*. 2014;9(3):e91340.
22. Gray J, Millett C, O'Sullivan C, Omar RZ, Majeed A. Association of age, sex and deprivation with quality indicators for diabetes: population-based cross sectional survey in primary care. *Journal of the Royal Society of Medicine*. 2006;99(11):576-81.
23. Zhao M, Woodward M, Vaartjes I, Millett ERC, Klipstein-Grobusch K, Hyun K, et al. Sex Differences in Cardiovascular Medication Prescription in Primary Care: A Systematic Review and Meta-Analysis. *J Am Heart Assoc*. 2020;9(11):e014742.
24. Zhao M, Vaartjes I, Graham I, Grobbee D, Spiering W, Klipstein-Grobusch K, et al. Sex differences in risk factor management of coronary heart disease across three regions. *Heart*. 2017;103(20):1587-94.
25. De Smedt D, De Bacquer D, De Sutter J, Dallongeville J, Gevaert S, De Backer G, et al. The gender gap in risk factor control: Effects of age and education on the control of cardiovascular risk factors in male and female coronary patients. The EUROASPIRE IV study by the European Society of Cardiology. *International Journal of Cardiology*. 2016;209:284-90.
26. Zhao M, Vaartjes I, Graham I, Grobbee D, Spiering W, Klipstein-Grobusch K, et al. Sex differences in risk factor management of coronary heart disease across three regions. *Heart*. 2017;103(20):1587.
27. Simpson CR, Hannaford PC, Williams D. Evidence for inequalities in the management of coronary heart disease in Scotland. *Heart (British Cardiac Society)*. 2005;91(5):630-4.
28. Chew DP, Juergens C, French J, Parsonage W, Horsfall M, Brieger D, et al. An examination of clinical intuition in risk assessment among acute coronary syndromes patients: observations from a prospective multi-center international observational registry. *Int J Cardiol*. 2014;171(2):209-16.
29. Mosca L, Mochari-Greenberger H, Dolor RJ, Newby LK, Robb KJ. Twelve-year follow-up of American women's awareness of cardiovascular disease risk and barriers to heart health. *Circ Cardiovasc Qual Outcomes*. 2010;3(2):120-7.
30. Lee RRS, Samsudin MI, Thirumoorthy T, Low LL, Kwan YH. Factors affecting follow-up non-attendance in patients with Type 2 diabetes mellitus and hypertension: a systematic review. *Singapore Med J*. 2019;60(5):216-23.
31. de Jong M, Vos RC, de Ritter R, van der Kallen CJ, Sep SJ, Woodward M, et al. Sex differences in cardiovascular risk management for people with diabetes in primary care: a cross-sectional study. *BJGP Open*. 2019;3(2):bjgpopen19X101645.
32. González-Pérez A, Saez M, Vizcaya D, Lind M, Garcia Rodriguez L. Incidence and risk factors for mortality and end-stage renal disease in people with type 2 diabetes and diabetic kidney disease: a population-based cohort study in the UK. *BMJ Open Diabetes Res Care*. 2021;9(1).
33. Molokhia M, Okoli GN, Redmond P, Asgari E, Shaw C, Schofield P, et al. Uncoded chronic kidney disease in primary care: a cross-sectional study of inequalities and cardiovascular disease risk management. *Br J Gen Pract*. 2020;70(700):e785-e92.
34. Kurani SS, Lampman MA, Funni SA, Giblon RE, Inselman JW, Shah ND, et al. Association Between Area-Level Socioeconomic Deprivation and Diabetes Care Quality in US Primary Care Practices. *JAMA Network Open*. 2021;4(12):e2138438-e.
35. Shippee ND, Shah ND, May CR, Mair FS, Montori VM. Cumulative complexity: a functional, patient-centered model of patient complexity can improve research and practice. *J Clin Epidemiol*. 2012;65(10):1041-51.

36. Ellis DA, McQueenie R, McConnachie A, Wilson P, Williamson AE. Demographic and practice factors predicting repeated non-attendance in primary care: a national retrospective cohort analysis. *Lancet Public Health*. 2017;2(12):e551-e9.
37. Chan JCN, Lim L-L, Wareham NJ, Shaw JE, Orchard TJ, Zhang P, et al. The Lancet Commission on diabetes: using data to transform diabetes care and patient lives. *The Lancet*. 2020;396(10267):2019-82.






figure 1a.eps

The percentage of patients with diabetic kidney disease monitored and managed in accordance with guidelines by sex, 2018



The percentage of patients with diabetic kidney disease monitored and managed in accordance with guidelines by ethnicity, 2018






figure 1d.eps

Category	Sub-category	Cost
All hospital visits (n=100)		
Hospital		
Medicine		400,000
Nursing		300,000
Outpatient		
Medicine		150,000
Nursing		150,000
Total		
		1,000,000

figure 2a.eps




figure 2d.eps

The image shows a screenshot of a table with the following content:

Table 3. Mean (SD) and 95% CI for variables per group	
Demographics	
Age	66.0 (4.4)
Sex	
Male	150 (45.5)
Female	175 (54.5)
Marital status	
Married	145 (44.5)
Single	180 (55.5)
Medical history	
Hypertension	105 (32.5)
Diabetes	120 (37.5)
Dyslipidemia	135 (41.5)
Atrial fibrillation	15 (4.7)
Stroke	25 (7.7)
Heart failure	35 (10.9)
Chronic kidney disease	45 (13.9)
COPD	55 (16.9)
Anemia	65 (20.0)
Medication	100 (30.9)
Functional status	
ADL	15 (4.7)
IADL	25 (7.7)
MDS	35 (10.9)
SPPB	45 (13.9)
Timed up and go	55 (16.9)
Gait speed	65 (20.0)
Handgrip strength	75 (23.0)
Muscle mass	85 (26.0)

figure 3c.eps

